Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

215. Proposed by **EDWIN L. RICH**, Lehigh University.

Solve (1)..........$x/a+y/b+c/z=3$,
(2)..........$x/a+b/y+z/c=3$,
(3)..........$a/x+y/b+z/c=3$.

216. Proposed by **L. E. NEWCOMB**, Los Gatos, Cal.

Express by radicals the roots of $x^6+ax^4+bx^3+\frac{1}{4}a^2x^2+\frac{1}{2}abx+c=0$.

217. Proposed by **F. P. MATZ**, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

Find the condition that $E\equiv x^5-bx^3+cx^2+dx-e$ shall be the product of a complete square and a complete cube.

218. Proposed by **SAUL EPSTEEN**.

Prove that $\sum_{r=0}^{n}\frac{c_r}{r+1}=\frac{2^{n+1}-1}{n+1}$ where c_r is the coefficient of x^r in the expansion of $(1+x)^n$.

GEOMETRY.

236. *Errata*. This problem should read: If two sides of a triangle pass through fixed points, the third side touches a fixed circle.

248. Proposed by **F. P. MATZ**, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

What is the equation to the curve on which lie the centers of the inscribed circles in the right-angled triangles of hypotenuse h?

244. Proposed by **O. W. ANTHONY**, Head of the Mathematical Department, DeWitt Clinton High School, New York.

Upon the sides of a triangle as bases isosceles triangles with base angles of 30° are constructed. Show that the lines joining the vertices of these isosceles triangles form an equilateral triangle.

245. Proposed by **W. J. GREENSTREET**, A. M., Editor of The Mathematical Gazette, Stroud, England.

PCP', DCD' are conjugate diameters of an ellipse; PN, DM are the ordinates to the major axis at P and D; show $CM/PN=CN/DM=AC/BC$, and that AP and BD' are parallel, and that AP' is parallel to BD.